

## Brilliant Violet 421™ anti-human CD107a (LAMP-1) Antibody

<b>Catalog# / Size</b>	328625 / 25 tests 328626 / 100 tests
<b>Clone</b>	H4A3
<b>Regulatory Status</b>	RUO
<b>Workshop</b>	P PR-63; BP 473; P P008
<b>Other Names</b>	Lysosome-Associated Membrane Protein 1, LGP-120, LAMP-1
<b>Isotype</b>	Mouse IgG1, κ
<b>Description</b>	CD107a, also known as Lysosome-Associated Membrane Protein 1 (LAMP-1) or LGP-120, is a 110-140 kD type I membrane glycoprotein. Mature CD107a is heavily glycosylated from a 40 kD core protein. This molecule is located on the luminal side of lysosomes. Upon activation, CD107a is transferred to the cell membrane surface of activated platelets, activated lymphocytes, macrophages, epithelial cells, endothelial cells, and some tumor cells. CD107a has been suggested to play a role in the protection of lysosomal membrane from lysosomal hydrolases which is involved in cell adhesion and regulation of tumor metastasis, and mediates autoimmune disease progression. CD107a is a ligand for galactin and E-selectin. Surface expression of LAMP-1 has been shown to correlate with CD8 <sup>+</sup> T cell and NK cell cytotoxicity.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Reported Reactivity</b>	African Green, Baboon, Chimpanzee, Cynomolgus, Pigtailed Macaque, Rhesus
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	Human adult adherent peripheral blood cells
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.  Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.  <a href="#">Learn more about Brilliant Violet™.</a>  This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.
<b>Excitation Laser</b>	Violet Laser (405 nm)

**Application Notes** Additional reported applications (for the relevant formats) include: Western blotting<sup>8</sup>, immunohistochemical staining<sup>2</sup>, immunofluorescence<sup>5,7</sup>, and immunoprecipitation<sup>5</sup>.

This antibody is specific to human LAMP-1. Positive control: Hela cells; LAMP-1 molecular weight appears to be at ~110 kDa on the gel due to high glycosylation.

#### Application References

(PubMed link indicates BioLegend citation)

1. Misse D, *et al.* 1999. *Blood* 93:2454.
2. Furuta K, *et al.* 2001. *Am. J. Pathol.* 159:449. (IHC)
3. Watanabe A, *et al.* 2011. *J. Biol. Chem.* 286:10702. [PubMed](#)
4. Baron Gaillard CL, *et al.* 2011. *Mol. Cell. Biol.* 22:5459. [PubMed](#)
5. Hauck CR and Meyer TF. 1997. *FEBS Lett.* 405:86. (IF, IP)
6. De Keersmaecker B, *et al.* 2012. *J. Virol.* 86:9351. [PubMed](#)
7. Knodler LA, *et al.* 2010. *P. Natl. Acad. Sci. USA.* 107:17733. (IF)
8. Oh J, *et al.* 2000. *Hum. Mol. Genet.* 9:375. (WB)
9. Salio M, *et al.* 2013 *PNAS.* 110:4753. [PubMed](#)

#### Product Citations

1. Keersmaecker B, *et al.* 2012. *J Virol. J. Virol.* [PubMed](#)
2. Braun M, *et al.* 2014. *PLoS Pathog.* 10:1004521. [PubMed](#)
3. Rider P, *et al.* 2015. *J Immunol.* 195: 1705-1712. [PubMed](#)
4. Terawaki S, *et al.* 2015. *J Cell Biol.* 210: 1133-1152. [PubMed](#)
5. Chalan P, *et al.* 2016. *J Rheumatol.* 43: 1008 - 1016. [PubMed](#)
6. Roberts E, *et al.* 2016. *PLoS One.* 11:e0168488. [PubMed](#)
7. Clayton KL, *et al.* 2021. *Cell Host Microbe.* 29(3):435-447.e9. [PubMed](#)
8. Motozono C, *et al.* 2021. *Cell Host Microbe.* . [PubMed](#)
9. Jackson E, *et al.* 2017. *PLoS One.* 10.1371/journal.pone.0185160. [PubMed](#)
10. Sagebiel AF, *et al.* 2019. *Nat Commun.* 10:975. [PubMed](#)
11. Sim MJW, *et al.* 2019. *Proc Natl Acad Sci U S A.* 116:12964. [PubMed](#)
12. Dai HS *et al.* 2017. *Immunity.* 47(1):159-170 . [PubMed](#)
13. Körner C *et al.* 2017. *Cell host & microbe.* 22(1):111-119 . [PubMed](#)

#### RRID

AB\_10899581 (BioLegend Cat. No. 328625)  
AB\_11203537 (BioLegend Cat. No. 328626)

## Antigen Details

---

<b>Structure</b>	LAMP-1 is a 417 amino acid protein with a molecular mass of 45 kD.
<b>Distribution</b>	Macrophages, epithelial cells, endothelial cells, some tumor cells; located on the luminal side of lysosomes or on the surface of cell membranes
<b>Function</b>	Protect lysosomal membrane from lysosomal hydrolases, adhesion
<b>Ligand/Receptor</b>	Galaptin
<b>Cell Type</b>	Endothelial cells, Epithelial cells, Macrophages
<b>Biology Area</b>	Cell Biology, Immunology, Neurodegeneration, Neuroscience, Protein Trafficking and Clearance
<b>Molecular Family</b>	Adhesion Molecules, CD Molecules
<b>Antigen References</b>	<ol style="list-style-type: none"><li>1. Sarafian V, <i>et al.</i> 2006. <i>Arch. Dermatol. Res.</i> 298:7381.</li><li>2. Schlossman SF, <i>et al.</i> 1995. <i>Leukocyte Typing V:White Cell Differentiation Antigens.</i> New York:Oxford University Press.</li><li>3. Sawada R, <i>et al.</i> 1993. <i>J. Biol. Chem.</i> 268:12675.</li><li>4. Chen JW, <i>et al.</i> 1988. <i>J. Biol. Chem.</i> 263:8754.</li><li>5. Chen JW, <i>et al.</i> 1986. <i>Biochem. Soc. Symp.</i> 51:97112.</li></ol>
<b>Gene ID</b>	<a href="#">3916</a>

## Related Protocols

---

[Cell Surface Flow Cytometry Staining Protocol](#)

## Other Formats


---

Biotin anti-human CD107a (LAMP-1), Purified anti-human CD107a (LAMP-1), FITC anti-human CD107a (LAMP-1), PE anti-human

CD107a (LAMP-1), Alexa Fluor® 488 anti-human CD107a (LAMP-1), Alexa Fluor® 647 anti-human CD107a (LAMP-1), PerCP/Cyanine5.5 anti-human CD107a (LAMP-1), APC anti-human CD107a (LAMP-1), Pacific Blue™ anti-human CD107a (LAMP-1), Brilliant Violet 421™ anti-human CD107a (LAMP-1), PE/Cyanine7 anti-human CD107a (LAMP-1), APC/Cyanine7 anti-human CD107a (LAMP-1), Brilliant Violet 510™ anti-human CD107a (LAMP-1), Brilliant Violet 605™ anti-human CD107a (LAMP-1), Purified anti-human CD107a (LAMP-1) (Maxpar® Ready), Brilliant Violet 650™ anti-human CD107a (LAMP-1), Brilliant Violet 711™ anti-human CD107a (LAMP-1), PerCP anti-human CD107a (LAMP-1), Brilliant Violet 785™ anti-human CD107a (LAMP-1), PE/Dazzle™ 594 anti-human CD107a (LAMP-1), TotalSeq™-A0155 anti-human CD107a (LAMP-1), TotalSeq™-C0155 anti-human CD107a (LAMP-1), TotalSeq™-B0155 anti-human CD107a (LAMP-1), APC/Fire™ 750 anti-human CD107a (LAMP-1) Antibody, PE/Cyanine5 anti-human CD107a (LAMP-1)

## Product Data

---



Thrombin-activated human peripheral blood platelets were stained with CD107a (clone H4A3) Brilliant Violet 421™ (filled histogram) or mouse IgG1,k Brilliant Violet 421™ isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587