

Alexa Fluor® 647 anti-human IFN-γ Antibody

Catalog# / Size	502516 / 100 tests
Clone	4S.B3
Regulatory Status	RUO
Other Names	Interferon-γ, Immune interferon, Type II interferon, T cell interferon, Macrophage-activating factor (MAF), IFN-g, IFN-gamma
Isotype	Mouse IgG1, κ
Description	Interferon-γ is a potent multifunctional cytokine which is secreted primarily by activated NK cells and T cells. Originally characterized based on anti-viral activities, IFN-γ also exerts anti-proliferative, immunoregulatory, and proinflammatory activities. IFN-γ can upregulate MHC class I and II antigen expression by antigen-presenting cells.

Product Details

Verified Reactivity	Human
Reported Reactivity	Chimpanzee, Baboon, Cynomolgus, Rhesus
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Partially purified, native human IFN-γ
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with Alexa Fluor® 647 under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	ICFC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 μl per 10 ⁶ cells in 100 μl volume. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

Alexa Fluor® and Pacific Blue™ are trademarks of Life Technologies Corporation.

[View full statement regarding label licenses](#)

Excitation Laser Red Laser (633 nm)

Application Notes

ELISA or ELISPOT Detection⁵: The biotinylated 4S.B3 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with purified NIB42 antibody (Cat. No. 502402/502404) or purified MD-1 antibody (Cat. No. 507502/507513) as the capture antibody.

Flow Cytometry^{3,4,6-8}: The fluorochrome-labeled 4S.B3 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IFN-γ -producing cells within mixed cell populations.

Additional reported applications (for the relevant formats) include: neutralization^{1,2}, Western blotting, immunohistochemical staining of paraformaldehyde-fixed, saponin-treated tissue sections, and immunocytochemistry. The 4S.B3 antibody can neutralize the bioactivity of natural or recombinant IFN-γ.

Note: For testing human IFN-γ in serum or plasma, BioLegend's ELISA Max™ Sets (Cat. No.

430101 to 430106) are specially developed and recommended.

Application References

(PubMed link indicates BioLegend citation)

1. Meager A, *et al.* 1984. *J. Interferon Res.* 4:619. (Neut)
2. Meager A, 1987. *Lymphokines and Interferons: A Practical Approach.* IRL Press Ltd, Oxford, p. 105. (Neut)
3. Sester M, *et al.* 2002. *J. Virol.* 76:3748. (ICFC)
4. Infante-Duarte C, *et al.* 2000 *J. Immunol.* 165:6107. (ICFC)
5. Goodier M, *et al.* 2000. *J. Immunol.* 165:139. (ELISA)
6. Chen H, *et al.* 2005. *J. Immunol.* 175:591. (ICFC)
7. Smeltz RB, 2007. *J. Immunol.* 178:4786. (ICFC)
8. Iwamoto S, *et al.* 2007. *J. Immunol.* 179:1449. (ICFC) [PubMed](#)
9. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (ICFC)

Product Citations

1. Yin S, *et al.* 2015. *Sci Rep.* 5: 14432. [PubMed](#)
2. Lever M, *et al.* 2016. *Proc Natl Acad Sci U S A.* 113: E6630 - E6638. [PubMed](#)
3. Corrado M, *et al.* 2020. *Cell Metab.* 32:981. [PubMed](#)
4. Hoang TN, *et al.* 2021. *Cell.* 184:460. [PubMed](#)
5. Oh DY, *et al.* 2020. *Cell.* 181:1612. [PubMed](#)
6. Wang Z, *et al.* 2020. *J Clin Invest.* 130:3717. [PubMed](#)
7. Bradley T *et al.* 2018. *Cell.* 175(2):387-399. [PubMed](#)
8. Li W, *et al.* 2020. *Immunity.* 53(2):456-470. [PubMed](#)

RRID

AB_493031 (BioLegend Cat. No. 502516)

Antigen Details

Structure	Cytokine; dimer; 20-25 kD (Mammalian)
Bioactivity	Antiviral/antiparasitic activities; inhibits proliferation; enhances MHC class I and II expression on APC
Cell Sources	CD8 ⁺ and CD4 ⁺ T cells, NK cells
Cell Targets	T cells, B cells, macrophages, NK cells, endothelial cells, fibroblasts
Receptors	IFN-γRα (CDw119) dimerized with IFN-γRβ (AF-1)
Cell Type	Tregs
Biology Area	Cell Biology, Immunology, Neuroinflammation, Neuroscience
Molecular Family	Cytokines/Chemokines
Antigen References	<ol style="list-style-type: none">1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook.</i> Academic Press, San Diego.2. De Maeyer E, <i>et al.</i> 1992. <i>Curr. Opin. Immunol.</i> 4:321.3. Farrar M, <i>et al.</i> 1993. <i>Annu. Rev. Immunol.</i> 11:571.4. Gray P, <i>et al.</i> 1987. <i>Lymphokines</i> 13:151.
Regulation	Upregulated by IL-2, FGF-basic, EGF; downregulated by vitamin D3 or DMN; labile at pH2
Gene ID	3458

Related Protocols


[Intracellular Cytokine Staining Protocol - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)


Other Formats

PE anti-human IFN-γ, APC anti-human IFN-γ, FITC anti-human IFN-γ, Biotin anti-human IFN-γ, Purified anti-human IFN-γ, Alexa Fluor® 488 anti-human IFN-γ, Alexa Fluor® 647 anti-human IFN-γ, Alexa Fluor® 700 anti-human IFN-γ, Pacific Blue™ anti-human IFN-γ, PerCP/Cyanine5.5 anti-human IFN-γ, APC/Cyanine7 anti-human IFN-γ, PE/Cyanine7 anti-human IFN-γ, Brilliant Violet 421™ anti-human IFN-γ, Brilliant Violet 570™ anti-human IFN-γ, Brilliant Violet 605™ anti-human IFN-γ, Brilliant Violet 650™ anti-human IFN-γ, Brilliant Violet 711™ anti-human IFN-γ, Brilliant Violet 785™ anti-human IFN-γ, Brilliant Violet 510™ anti-human IFN-γ, PE/Dazzle™ 594 anti-human IFN-γ, APC/Fire™ 750 anti-human IFN-γ, PerCP anti-human IFN-γ, Brilliant Violet 750™ anti-human

Product Data



PMA/ionomycin-stimulated (6 hours)
human peripheral blood lymphocytes
stained with mouse IgG1 Alexa Fluor®
647 isotype control and CD3 (UCHT1)
PE



PMA/ionomycin-stimulated (6 hours)
human peripheral blood lymphocytes
stained with 4S.B3 Alexa Fluor® 647 and
CD3 (UCHT1) PE

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587