

## TotalSeq™-B0050 anti-human CD19 Antibody

<b>Catalog# / Size</b>	302263 / 10 µg
<b>Clone</b>	HIB19
<b>Regulatory Status</b>	RUO
<b>Workshop</b>	V CD19.11
<b>Other Names</b>	B4
<b>Isotype</b>	Mouse IgG1, κ
<b>Barcode Sequence</b>	CTGGGCAATTACTCG
<b>Description</b>	CD19 is a 95 kD type I transmembrane glycoprotein also known as B4. It is a member of the immunoglobulin superfamily expressed on B-cells (from pro-B to blastoid B cells, absent on plasma cells) and follicular dendritic cells. CD19 is involved in B cell development, activation, and differentiation. CD19 forms a complex with CD21 (CR2) and CD81 (TAPA-1), and functions as a BCR co-receptor.

### Product Details

---

<b>Verified Reactivity</b>	Human, Cynomolgus, Rhesus
<b>Reported Reactivity</b>	Chimpanzee
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 1 mM EDTA.
<b>Preparation</b>	The antibody was purified by chromatography and conjugated with TotalSeq™-B oligomer under optimal conditions.
<b>Concentration</b>	0.5 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">PG - Quality tested</a>
<b>Recommended Usage</b>	<p>Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> and the oligomer sequence is confirmed by sequencing. TotalSeq™-B antibodies are compatible with 10x Genomics Single Cell Gene Expression <a href="#">Solutions</a>.</p> <p>To maximize performance, it is strongly recommended that the reagent be titrated for each application, and that you centrifuge the antibody dilution before adding to the cells at 14,000xg at 2 - 8°C for 10 minutes. Carefully pipette out the liquid avoiding the bottom of the tube and add to the cell suspension. For Proteogenomics analysis, the suggested starting amount of this reagent for titration is ≤ 1.0 µg per million cells in 100 µL volume. Refer to the corresponding TotalSeq™ protocol for specific staining instructions.</p> <p>Buyer is solely responsible for determining whether Buyer has all intellectual property rights that are necessary for Buyer's intended uses of the BioLegend TotalSeq™ products. For example, for any technology platform Buyer uses with TotalSeq™, it is Buyer's sole responsibility to determine whether it has all necessary third party intellectual property rights to use that platform and TotalSeq™ with that platform.</p>
<b>Application Notes</b>	<p>Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections<sup>8</sup> and blocking of B cell proliferation. Clone HIB19 is not recommended for formalin-fixed paraffin-embedded sections. The Ultra-LEAF™ purified antibody (Endotoxin &lt; 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 302267 &amp; 302268).</p> <p>Clone HIB19 partially blocks anti-human CD19 clones 4G7 and SJ25C1 staining based on in-house testing</p>

**Additional Product Notes** TotalSeq™ reagents are designed to profile protein levels at a single cell level following an optimized protocol similar to the CITE-seq workflow. A compatible single cell device (e.g. [10x Genomics Chromium System and Reagents](#)) and sequencer (e.g. Illumina analyzers) are required. Please contact [technical support](#) for more information, or visit [biolegend.com/totalseq](#).

The barcode flanking sequences are GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTNNNNNNNNNN (PCR handle), and NNNNNNNNGCTTTAAGGCCGGTCCTAGC\*A\*A (capture sequence). N represents either randomly selected A, C, G, or T, and \* indicates a phosphorothioated bond, to prevent nuclease degradation.

View more applications data for this product in our [Scientific Poster Library](#).

## Application References

(PubMed link indicates BioLegend citation)

1. Schlossman S, *et al.* 1995. Leucocyte Typing V. Oxford University Press. New York.
2. Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
3. Bradbury L, *et al.* 1993. *J. Immunol.* 151:2915.
4. Joseph A, *et al.* 2010. *J. Virol.* 84:6645. [PubMed](#)
5. Wang X, *et al.* 2010. *Haematologica.* 95:884. (FC) [PubMed](#)
6. Walker JD, *et al.* 2009. *J. Immunol.* 182:1548. (Block) [PubMed](#)
7. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
8. Hansen A, *et al.* 2002. *Arthritis Rheum.* 46:2160. (IHC)
9. Stoeckius M, *et al.* 2017. *Nat. Methods.* 14:865. (PG)
10. Peterson VM, *et al.* 2017. *Nat. Biotechnol.* 35:936. (PG)

**RRID** AB\_2800740 (BioLegend Cat. No. 302263)

## Antigen Details

<b>Structure</b>	Ig superfamily, type I transmembrane glycoprotein, 95 kD
<b>Distribution</b>	B lineage (except plasma cells), follicular dendritic cells
<b>Function</b>	B cell activation and differentiation
<b>Ligand/Receptor</b>	Forms complex with CD21 (CR2) and CD81 (TAPA-1), BCR coreceptor
<b>Cell Type</b>	B cells, Dendritic cells
<b>Biology Area</b>	Costimulatory Molecules, Immunology
<b>Molecular Family</b>	CD Molecules
<b>Antigen References</b>	1. Tedder T, <i>et al.</i> 1994. <i>Immunol. Today</i> 15:437. 2. Bradbury L, <i>et al.</i> 1993. <i>J. Immunol.</i> 151:2915.
<b>Gene ID</b>	<a href="#">930</a>

## Related Protocols

[TotalSeq™-B or -C with 10x Feature Barcoding Technology](#)

## Other Formats

APC anti-human CD19, Biotin anti-human CD19, FITC anti-human CD19, PE anti-human CD19, PE/Cyanine5 anti-human CD19, Purified anti-human CD19, APC/Cyanine7 anti-human CD19, PE/Cyanine7 anti-human CD19, Alexa Fluor® 488 anti-human CD19, Alexa Fluor® 647 anti-human CD19, Pacific Blue™ anti-human CD19, Alexa Fluor® 700 anti-human CD19, PerCP anti-human CD19, PerCP/Cyanine5.5 anti-human CD19, Brilliant Violet 421™ anti-human CD19, Brilliant Violet 570™ anti-human CD19, Brilliant Violet 650™ anti-human CD19, Brilliant Violet 785™ anti-human CD19, Brilliant Violet 510™ anti-human CD19, Brilliant Violet 605™ anti-human CD19, Brilliant Violet 711™ anti-human CD19, Purified anti-human CD19 (Maxpar® Ready), Alexa Fluor® 594 anti-human CD19, PE/Dazzle™ 594 anti-human CD19, APC/Fire™ 750 anti-human CD19, TotalSeq™-A0050 anti-human CD19, Brilliant Violet 750™ anti-human CD19, TotalSeq™-B0050 anti-human CD19, TotalSeq™-C0050 anti-human CD19, Spark NIR™ 685 anti-human CD19, Ultra-LEAF™ Purified anti-human CD19, APC/Fire™ 810 anti-human CD19, PE/Fire™ 640 anti-human CD19, PE/Fire™ 700 anti-human CD19, TotalSeq™-D0050 anti-human CD19, Spark YG™ 593 anti-human CD19, GMP Pacific Blue™ anti-human CD19, Spark Violet™ 423 anti-human CD19, GMP PE anti-human CD19, GMP APC anti-human CD19, KIRAVIA Blue 520™ anti-human CD19, GMP PerCP/Cyanine5.5 anti-human CD19, GMP PE/Cyanine7 anti-human CD19

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587