

Recombinant Human IL-6 (carrier-free)

Catalog# / Size	570802 / 10 µg 570804 / 25 µg 570806 / 100 µg 570808 / 500 µg
Regulatory Status	RUO
Other Names	Interleukin-6, B cell stimulating factor-2 (BSF-2), Cytotoxic T cell differentiation factor (CDF), Hepatocyte stimulating factor (HSF), Hybridoma/plasmacytoma growth factor (HPGF)
Description	IL-6 is a potent lymphoid cell growth factor that stimulates the growth and survival of certain B cells and T cells. IL-6 plays a role in host defense, acute phase reactions, immune response, and hematopoiesis. IL-6 is expressed by T cells, B cells, monocytes, fibroblasts, hepatocytes, endothelial cells and keratinocytes.

Product Details

Source	Human IL-6, amino acids Pro29-Met212 (Accession # NM_000600) was expressed in <i>E. coli</i> .
Molecular Mass	The 185 amino acid N-terminal methionylated recombinant protein has a predicted molecular mass of 21,041 Da. The DTT-reduced and non-reduced protein migrate at approximately 23kDa by SDS-PAGE.
Purity	Purity is >98%, as determined by Coomassie stained SDS-PAGE.
Formulation	0.22 µm filtered protein solution is in 10mM NaH ₂ PO ₄ , 150mM NaCl, pH 7.2.
Endotoxin Level	Endotoxin level is <0.1 EU/µg (<0.01ng/µg) protein as determined by the LAL method.
Concentration	10 and 25 µg sizes are bottled at 200 µg/mL. 100 µg size and larger sizes are lot-specific and bottled at the concentration indicated on the vial. To obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.
Storage & Handling	Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to six months, or at -70°C or colder until the expiration date. For maximum results, quick spin vial prior to opening. The protein can be aliquoted and stored at -20°C or colder. Stock solutions can also be prepared at 50 - 100 µg/mL in appropriate sterile buffer, carrier protein such as 0.2 - 1% BSA or HSA can be added when preparing the stock solution. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. Avoid repeated freeze/thaw cycles.
Activity	<p>The ED₅₀ is 4-20 pg/ml, corresponding to a specific activity of 0.5-2.5 X10⁸ units/mg, as determined by a dose dependent stimulation in a 7TD1 cell proliferation assay.</p> <p>The specific activity of recombinant human IL-6 is approximately 5.0 x 10⁴ IU/µg when compared against the 1st WHO International Standard for Interleukin-6 (NIBSC code: 89/548) as determined by the dose dependent stimulation of 7TD1 cell proliferation.</p> <p>For more information on specific activity, please visit the Recombinant Protein Unit Conversions page.</p>
Application	Bioassay
Recommended Usage	Use when high specific biological activity is required.
Application Notes	BioLegend carrier-free recombinant proteins provided in liquid format are shipped on blue-ice. Our comparison testing data indicates that when handled and stored as recommended, the liquid format has equal or better stability and shelf-life compared to commercially available lyophilized proteins after reconstitution. Our liquid proteins are verified in-house to maintain activity after shipping on blue ice and are backed by our 100% satisfaction guarantee . If you have any concerns, contact us at tech@biolegend.com .
Additional Product Notes	View more applications data for this product in our Scientific Poster Library .
Application References	

1. Reitsma JM, *et al.* 2013. *J Virol.* 87:10763. [PubMed](#)


Product Citations

1. Ibrahim ML, *et al.* 2018. *Cell Rep.* 25:3036. [PubMed](#)
2. Fulciniti M, *et al.* 2018. *Cell Rep.* 25:3693. [PubMed](#)
3. Wong GK, *et al.* 2019. *JCI Insight.* 4. [PubMed](#)
4. Jokela TA, *et al.* 2018. *Front Cell Dev Biol.* 6:41. [PubMed](#)
5. Zhang H, *et al.* 2020. *Front Cell Dev Biol.* 8:205. [PubMed](#)
6. Guo J, *et al.* 2019. *Cancer Immunol Res.* 1.349305556. [PubMed](#)
7. Reitsma J, *et al.* 2013. *J Virol.* 87:10763. [PubMed](#)
8. Sugawara E, *et al.* 2020. *Autophagy.* 1.323611111. [PubMed](#)

Antigen Details

Structure	Neuropoietic cytokine family
Distribution	T cells, B cells, macrophages, bone marrow stromal cells, fibroblasts, keratinocytes, mesangium, astrocytes, endothelial cells
Function	Upregulated by IL-1, TNF, PDGF, IFN- β , TNF- α , NGF, IL-17; downregulated by glucocorticoids IL-4, TGF- β
Interaction	T cells, B cells, hepatocytes, cholinergic neurons
Ligand/Receptor	Heterodimer IL-6R α (CD126)/IL-6R β (CD130, gp130)
Bioactivity	Hematopoiesis; antigen-specific immune responses, inflammatory reactions; acute phase reaction
Cell Type	Hematopoietic stem and progenitors
Biology Area	Cell Biology, Immunology, Innate Immunity, Stem Cells
Molecular Family	Cytokines/Chemokines
Antigen References	<ol style="list-style-type: none">1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook.</i> Academic Press San Diego.2. Hirano T. 1998. <i>Intl. Rev. Immunol.</i> 16:249.3. Patterson P. 1992. <i>Curr. Opin. Neurobiol.</i> 2:94.4. Van Oers M, <i>et al.</i> 1993. <i>Ann. Hematology</i> 66:219.
Gene ID	3569

Product Data



Recombinant human IL-6 induces the proliferation of 7TD1 cells in a dose-dependent manner. BioLegend's product was compared side-by-side to the leading competitor equivalent product.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587