

PE anti-human IL-1RL1 (ST2) Antibody

Catalog# / Size	379403 / 25 tests 379404 / 100 tests
Clone	A16008J
Regulatory Status	RUO
Other Names	IL-33R, ST2, IL-1RL1
Isotype	Mouse IgG2b, κ
Description	IL1RL1 is a member of the Toll-like receptor superfamily the receptor for Interleukin-33. It is a 556 amino acid single-pass type I membrane protein. ST2 exists in two forms as splice variants: a soluble form, which acts as a decoy receptor and sequesters free IL-33, and a membrane-bound form (ST2), which activates the MyD88/NF-κB signaling pathway. Signaling through IL1RL1 drives cytokine production in mast cells, basophils, eosinophils, invariant NKT and NK cells, Th2 lymphocytes, and type 2 innate immune cells. IL-33 signaling is required for maintaining the suppressive function of regulatory T cells.

Product Details

Verified Reactivity	Human
Antibody Type	Monoclonal
Host Species	Mouse
Immunogen	Recombinant human IL-1RL1, AA19-328
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
Preparation	The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions.
Concentration	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our Concentration and Expiration Lookup or Certificate of Analysis online tools.)
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Application	FC - Quality tested
Recommended Usage	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis . For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Excitation Laser	Blue Laser (488 nm) Green Laser (532 nm)/Yellow-Green Laser (561 nm)
RRID	AB_2922608 (BioLegend Cat. No. 379403) AB_2922608 (BioLegend Cat. No. 379404)

Antigen Details

Structure	Single-pass type I membrane protein
Distribution	Th2 lymphocytes, mast cells, eosinophils, basophils, smooth muscle cells, endothelial cells
Function	Cytokine signaling
Interaction	Signaling requires association of the coreceptor IL1RAP. Recruits MYD88, IRAK1, IRAK4, and TRAF6.

Ligand/Receptor IL-33

Antigen References

1. Schmitz J, et al. 2005. *Immunity*. 23: 479-490.
2. Brint E, et al. 2004. *Nat Immun*. 5:373-379.
3. Griesenauer B, et al. 2017. *Front Immunol*. 8:475.

Gene ID [9173](#)


Related Protocols

[Cell Surface Flow Cytometry Staining Protocol](#)

Other Formats

Purified anti-human IL-1RL1 (ST2), PE anti-human IL-1RL1 (ST2)

Product Data



Human peripheral blood mononuclear cells were stimulated with IL-3 overnight and stained with anti-human HLA-DR APC, anti-human CD123 PE/Dazzle 594™ and anti-human IL-1RL1 (ST2) (clone A16008J) PE (left) or mouse IgG2b, κ PE isotype control (right).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587