

## PerCP/Cyanine5.5 anti-human CD27 Antibody

<b>Catalog# / Size</b>	356407 / 25 tests 356408 / 100 tests
<b>Clone</b>	M-T271
<b>Regulatory Status</b>	RUO
<b>Workshop</b>	V 5T CD27.03
<b>Other Names</b>	S152, T14, TNFRSF7
<b>Isotype</b>	Mouse IgG1, $\kappa$
<b>Description</b>	CD27 is a 50-55 kD type I membrane protein also known as S152 and T14. It is a lymphocyte-specific member of the TNF-receptor superfamily. CD27 is expressed on medullary thymocytes, virtually all mature T cells, some B cells, and NK cells. CD27 binds to CD70, and plays a role in costimulation of T cell activation and regulation of B cell differentiation and proliferation. The cytoplasmic domains of CD27 have also been shown to interact with TRAF2 and TRAF5 to elicit NF- $\kappa$ B and SAPK/JNK activation.

### Product Details

---

<b>Verified Reactivity</b>	Human
<b>Reported Reactivity</b>	Baboon, Cynomolgus, Rhesus
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Mouse
<b>Immunogen</b>	Human T cells from a T-ALL patient.
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions.
<b>Concentration</b>	Lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">FC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is 5 $\mu$ l per million cells in 100 $\mu$ l staining volume or 5 $\mu$ l per 100 $\mu$ l of whole blood.  * PerCP/Cyanine5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.
<b>Application Notes</b>	Additional reported applications (for the relevant formats) include: immunohistochemical staining of formalin-fixed paraffin-embedded frozen tissue sections <sup>1</sup> , immunofluorescent staining <sup>2</sup> , and ELISA <sup>3</sup> .
<b>Additional Product Notes</b>	BioLegend is in the process of converting the name PerCP/Cy5.5 to PerCP/Cyanine5.5. The dye molecule remains the same, so you should expect the same quality and performance from our PerCP/Cyanine5.5 products. Contact <a href="#">Technical Service</a> if you have any questions.
<b>Application References</b>	1. Ma S, <i>et al.</i> 2011. <i>J. Virol.</i> 85:165. (IHC) 2. Manzo A, <i>et al.</i> 2008. <i>Arthritis Rheum.</i> 11:3377. (IF) 3. Kato K, <i>et al.</i> 2007. <i>Exp. Hematol.</i> 35:434. (ELISA)
<b>(PubMed link indicates BioLegend citation)</b>	
<b>Product Citations</b>	1. Lindesmith LC, <i>et al.</i> 2020. <i>Cell Mol Gastroenterol Hepatol.</i> 0.586805556. <a href="#">PubMed</a> 2. Gurusamy D, <i>et al.</i> 2020. <i>Cancer Cell.</i> 37(6):818-833.e9. <a href="#">PubMed</a>

3. Sutton HJ, *et al.* 2021. Cell Reports. 34(6):108684. [PubMed](#)
4. Zumaquero E, *et al.* 2019. Elife. 8. [PubMed](#)
5. Li X, *et al.* 2019. Front Oncol. 1.115277778. [PubMed](#)
6. Upasani V, *et al.* 2019. Front Immunol. 2.152777778. [PubMed](#)

**RRID** AB\_2561905 (BioLegend Cat. No. 356407)  
AB\_2561906 (BioLegend Cat. No. 356408)

## Antigen Details

---

<b>Structure</b>	TNF-R superfamily, type I transmembrane glycoprotein, 50-55 kD
<b>Distribution</b>	Medullary thymocytes, T and B cell subsets, NK cells
<b>Interaction</b>	CD27 binds to CD70
<b>Ligand/Receptor</b>	CD70
<b>Cell Type</b>	B cells, NK cells, T cells, Thymocytes
<b>Biology Area</b>	Costimulatory Molecules, Immunology
<b>Molecular Family</b>	CD Molecules
<b>Antigen References</b>	<ol style="list-style-type: none"><li>1. Knapp W, <i>et al.</i> 1989. Leucocyte Typing IV: White Cell Differentiation Antigens. Oxford University Press.</li><li>2. Schlossman S, <i>et al.</i> 1995. Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press.</li><li>3. Hintzen R, <i>et al.</i> 1994. <i>Immunol. Today</i> 15:307.</li><li>4. Agematsu K, <i>et al.</i> 1995. <i>J. Immunol.</i> 154:3627.</li></ol>
<b>Gene ID</b>	<a href="#">939</a>

## Related Protocols

---

[Cell Surface Flow Cytometry Staining Protocol](#)


## Other Formats

---

Purified anti-human CD27, FITC anti-human CD27, PE anti-human CD27, PerCP/Cyanine5.5 anti-human CD27, APC anti-human CD27, PE/Cyanine7 anti-human CD27, Pacific Blue™ anti-human CD27, Alexa Fluor® 700 anti-human CD27, Brilliant Violet 421™ anti-human CD27, Brilliant Violet 510™ anti-human CD27, PE/Dazzle™ 594 anti-human CD27, APC/Cyanine7 anti-human CD27, Biotin anti-human CD27, APC/Fire™ 750 anti-human CD27, Brilliant Violet 711™ anti-human CD27, PerCP anti-human CD27, Alexa Fluor® 647 anti-human CD27, KIRAVIA Blue 520™ anti-human CD27, PE/Cyanine5 anti-human CD27

## Product Data

---



Human peripheral blood lymphocytes were stained with CD27 (clone M-T271) PerCP/Cyanine5.5 (filled histogram), or mouse IgG1, ? PerCP/Cyanine5.5 isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587