

Purified anti-human IL-4 (Maxpar[®] Ready) Antibody

Catalog# / Size	500829 / 100 µg
Clone	MP4-25D2
Regulatory Status	RUO
Other Names	Interleukin-4, Ia inducing factor (IaIF), B-cell stimulating factor-1 (BSF-1), Hodgkin's cell growth factor (HCGF), Mast cell growth factor-2 (MCGF-2), Macrophage fusion factor (MFF), T cell growth factor-2 (TCGF-2)
Isotype	Rat IgG1, κ
Description	IL-4 is a pleiotropic cytokine that is produced by activated T cells, mast cells, and basophils. IL-4 elicits many different biological responses but has two dominant functions. The first is regulating differentiation of naïve CD4 ⁺ T cell to the Th2 type. Th2 cells produce IL-4, IL-5, IL-10, and IL-13, which tend to favor a humoral immune response while suppressing a cell-mediated immune response controlled by Th1 cells. The second is regulating IgE and IgG1 production by B cells.

Product Details

Verified Reactivity	Human
Reported Reactivity	Pig, Rhesus
Antibody Type	Monoclonal
Host Species	Rat
Immunogen	CHO-expressed, recombinant human IL-4
Formulation	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and EDTA.
Preparation	The antibody was purified by affinity chromatography.
Concentration	1.0 mg/ml
Storage & Handling	The antibody solution should be stored undiluted between 2°C and 8°C.
Application	ELISA - Quality tested CyTOF[®] - Verified
Recommended Usage	This product is suitable for use with the Maxpar[®] Metal Labeling Kits . For metal labeling using Maxpar [®] Ready antibodies, proceed directly to the step to Partially Reduce the Antibody by adding 100 µl of Maxpar [®] Ready antibody to 100 µl of 4 mM TCEP-R in a 50 kDa filter and continue with the protocol. Always refer to the latest version of Maxpar [®] User Guide when conjugating Maxpar [®] Ready antibodies.
Application Notes	ELISA Detection^{1,3} or ELISPOT Detection^{4,5}: The biotinylated MP4-25D2 antibody is useful as a detection antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with purified 8D4-8 antibody (Cat. No. 500702/500707) as the capture antibody. Flow Cytometry^{6,9}: The fluorochrome-labeled MP4-25D2 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IL-4 -producing cells within mixed cell populations. Neutralization¹⁻³: The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for neutralization of human IL-4 bioactivity (Cat. No. 500815). The MP4-25D2 antibody can neutralize the bioactivity of natural or recombinant IL-4.
Additional Product Notes	Maxpar [®] is a registered trademark of Standard BioTools Inc.
Application References	<ol style="list-style-type: none"> Chretien I, <i>et al.</i> 1989. <i>J. Immunol. Methods</i> 117:67. (ELISA Detection, Neut) Ramanathan L, <i>et al.</i> 1993. <i>Biochem.</i> 32:3549. (Neut) Abrams J, <i>et al.</i> 1992. <i>Immunol. Rev.</i> 127:5. (ELISA Detection, Neut) Mahanty S, <i>et al.</i> 1992. <i>J. Immunol.</i> 148:3567. (ELISPOT Detection) Klinman D, <i>et al.</i> 1994. <i>Curr. Prot. Immunol.</i> John Wiley and Sons New York. Unit 6.19. (ELISPOT)

- Detection)
6. Prussin C, *et al.* 1995. *J. Immunol. Methods* 188:117. (ICFC)
 7. Raqib R, *et al.* 1995. *Infect. Immun.* 63:289.
 8. Andersson J, *et al.* 1994. *Immunology* 83:16.
 9. Iwamoto S, *et al.* 2007. *J. Immunol.* 179:1449. (ICFC) [PubMed](#)
 10. Kubota M, *et al.* 1997. *J. Immunol.* 158:5321.
 11. Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113. [PubMed](#)
 12. Kroneke MA, *et al.* 2012. *J. Immunol.* 188:3734. [PubMed](#)

RRID AB_2562840 (BioLegend Cat. No. 500829)

Antigen Details

Structure	Cytokine; 15-19 kD (Mammalian)
Bioactivity	Differentiation of naïve CD4 ⁺ T cells to the T _H 2 type, proliferation/differentiation of activated B cells, expression of class II MHC antigens, and of low affinity IgE receptors in resting B cells
Cell Sources	Mast cells, T cells, bone marrow stromal cells
Cell Targets	B cells, T cells, monocytes, endothelial cells, fibroblasts
Receptors	Heterodimer IL-4R α (CD124); γ -subunit (CD132) in common with IL-2R, IL-7R, IL-13R, IL-15R
Cell Type	Tregs
Biology Area	Cell Biology, Immunology, Neuroinflammation, Neuroscience
Molecular Family	Cytokines/Chemokines
Antigen References	<ol style="list-style-type: none"> 1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook</i>. Academic Press San Diego. 2. Boulay J, <i>et al.</i> 1992. <i>Curr. Opin. Immunol.</i> 4:294. 3. Dullens H, <i>et al.</i> 1991. <i>In vivo</i> 5:567. 4. Paul W. 1991. <i>Blood</i> 77:1859.
Regulation	Upregulated by IL-2, platelet activating factor; downregulated by TGF- β
Gene ID	3565


Related Protocols

[Sandwich ELISA Protocol](#)


Other Formats

APC anti-human IL-4, Biotin anti-human IL-4, FITC anti-human IL-4, PE anti-human IL-4, Purified anti-human IL-4, Alexa Fluor® 488 anti-human IL-4, Alexa Fluor® 647 anti-human IL-4, Brilliant Violet 421™ anti-human IL-4, PerCP/Cyanine5.5 anti-human IL-4, PE/Cyanine7 anti-human IL-4, Brilliant Violet 605™ anti-human IL-4, Purified anti-human IL-4 (Maxpar® Ready), PE/Dazzle™ 594 anti-human IL-4, APC/Cyanine7 anti-human IL-4, Brilliant Violet 510™ anti-human IL-4, Ultra-LEAF™ Purified anti-human IL-4

Product Data



Human Th2-polarized CD4⁺ T cells were incubated for 5 hours in media alone (bottom) or with PMA and Ionomycin (top) in the presence of monensin and brefeldin A. Cells were then fixed, permeabilized, and stained with 145Nd anti-CD4 (RPA-T4) and 142Nd anti-IL-4 (MP4-25D2). Data provided by DVS Sciences.



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587