

## PerCP/Cyanine5.5 anti-mouse IL-12/IL-23 p40 (monomer, dimer, heterodimer) Antibody

<b>Catalog# / Size</b>	505211 / 25 µg 505212 / 100 µg
<b>Clone</b>	C15.6
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	Interleukin-12 p40, Interleukin-23 p40, Cytotoxic lymphocyte maturation factor (CLMF), Natural killer cell stimulatory factor (NKSF), CTL maturation factor (TcMF), T-cell stimulating factor (TSF)
<b>Isotype</b>	Rat IgG1, κ
<b>Description</b>	The C15.6 antibody reacts with mouse IL-12 p40 subunit of the IL-12 p70 and IL-23 p40 subunit of the IL-23 p19/p40, as well as p40 monomer and homodimer, or heterodimer. The C15.6 antibody can not neutralize the bioactivity of natural or recombinant IL-12.

### Product Details

<b>Verified Reactivity</b>	Mouse, IL-12/IL-23 p40 subunit (monomer, homodimer and heterodimer IL-12 p35/p40 or IL-23 p19/p40)
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Rat
<b>Immunogen</b>	CHO-expressed, recombinant mouse IL-12 p70
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions.
<b>Concentration</b>	0.2 mg/ml
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">ICFC - Quality tested</a>
<b>Recommended Usage</b>	Each lot of this antibody is quality control tested by <a href="#">intracellular immunofluorescent staining with flow cytometric analysis</a> . For flow cytometric staining, the suggested use of this reagent is =0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.  * PerCP/Cyanine5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.
<b>Application Notes</b>	<b>ELISA or ELISPOT Capture</b> <sup>1,2,4,6,8,10</sup> : The purified C15.6 antibody is useful as the capture antibody in a sandwich ELISA or ELISPOT assay, when used in conjunction with the biotinylated C17.8 (Cat. No. 505302) antibody as the detecting antibody. This assay detects p40 as monomer, homodimer, or heterodimer complexed with p35. The LEAF™ purified antibody is suggested for ELISPOT capture. <b>Additional reported applications (for the relevant formats) include:</b> immunohistochemical staining <sup>9</sup> of paraformaldehyde-fixed saponin-treated frozen tissue sections, immunoprecipitation <sup>3</sup> , and Western blotting <sup>3</sup> . <b>Note:</b> For testing mouse IL-12 p40 (monomer, dimer, heteromer) in serum or plasma, BioLegend's ELISA Max™ Sets (Cat. No. 431601 to 431606) are specially developed and recommended.
<b>Additional Product Notes</b>	BioLegend is in the process of converting the name PerCP/Cy5.5 to PerCP/Cyanine5.5. The dye molecule remains the same, so you should expect the same quality and performance from our PerCP/Cyanine5.5 products. Contact <a href="#">Technical Service</a> if you have any questions.
<b>Application References</b>	1. Kitagaki K, <i>et al.</i> 2002. <i>Clin. Diagn. Lab Immunol.</i> 9:1260. 2. Reichmann G, <i>et al.</i> 1999. <i>J. Immunol.</i> 163:3354.

**(PubMed link indicates BioLegend citation)**

3. Wysocka M, *et al.* 1995. *Eur. J. Immunol.* 25:672.
4. Gately M. 1995. *Curr. Prot. Immunol.* John Wiley and Sons, New York. Unit 6.16.
5. Macatonia S, *et al.* 1995. *J. Immunol.* 154:5071.
6. O'Connell PJ, *et al.* 2006. *Blood* 107:1010.
7. Akilov OE, *et al.* 2007. *J. Leukocyte Biol.* 2007;10.1189/jlb.0706439.
8. Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113.
9. Huang LY, *et al.* 2001. *J. Immunol.* 167:1423.
10. Xu G, *et al.* 2007. *J. Immunol.* 179:5358. [PubMed](#)
11. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
12. Malu DT, *et al.* 2011. *J. Immunol.* 186:6271. [PubMed](#)

**Product Citations**

1. Weinstock NI, *et al.* 2020. *Neuron.* 107(1):65-81. [PubMed](#)

**RRID**

AB\_2566224 (BioLegend Cat. No. 505211)  
AB\_2566225 (BioLegend Cat. No. 505212)

## Antigen Details

---

<b>Structure</b>	Cytokine; monomer, heterodimer (p40:p35 or p40:p19) or homodimer (p40:p40)
<b>Bioactivity</b>	IL-12 p70 (p40:p35) induces IFN- $\gamma$ , TNF- $\alpha$ production in T and NK cells; costimulation of PBL proliferation; proliferation/differentiation of T $\beta$ 1 T lymphocytes. IL-23 (p40:p19) induces proliferation and production of IFN- $\gamma$ by human me
<b>Cell Sources</b>	Dendritic cells, monocytes/macrophages, B cells, T cells
<b>Cell Targets</b>	T cells, NK cells
<b>Receptors</b>	IL-12R $\beta$ 1 binds p40; dimeric with IL-12R $\beta$ 2 binds p35
<b>Biology Area</b>	Immunology, Innate Immunity
<b>Molecular Family</b>	Cytokines/Chemokines
<b>Antigen References</b>	<ol style="list-style-type: none"><li>1. Fitzgerald K, <i>et al.</i> Eds. 2001. <i>The Cytokine FactsBook.</i> Academic Press, San Diego.</li><li>2. Quesniaux V. 1992. <i>Research Immunol.</i> 143:385.</li><li>3. Trinchieri G, <i>et al.</i> 1992 <i>Prog. Growth Factor Res.</i> 4:355.</li><li>4. Trinchieri G, <i>et al.</i> 1993 <i>Immunol. Today.</i> 14:335.</li><li>5. Oppmann B, <i>et al.</i> 2000 <i>Immunity.</i> 13:715.</li><li>6. Aggarwal S, <i>et al.</i> 2003 <i>J. Biol. Chem.</i> 278:1910.</li><li>7. Parham C, <i>et al.</i> 2002 <i>J. Immunol.</i> 168:5699.</li><li>8. Belladonna ML, <i>et al.</i> 2002 <i>J. Immunol.</i> 168:5448.</li><li>9. Lankford, CS, <i>et al.</i> 2003 <i>J. Leukocyte Biol.</i> 73:49.</li></ol>
<b>Regulation</b>	Downregulated by IL-10; homodimeric p40 antagonistic to functional p70 heterodimer; p40 monomer has no function; p40 subunit in common with IL-23
<b>Gene ID</b>	<a href="#">16160</a>

## Related Protocols

---

[Intracellular Cytokine Staining Protocol - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)


## Other Formats

---

APC anti-mouse IL-12/IL-23 p40 (monomer, dimer, heterodimer), PE anti-mouse IL-12/IL-23 p40 (monomer, dimer, heterodimer), Purified anti-mouse IL-12/IL-23 p40 (monomer, dimer, heterodimer), PE/Cyanine7 anti-mouse IL-12/IL-23 p40 (monomer, dimer, heterodimer), PerCP/Cyanine5.5 anti-mouse IL-12/IL-23 p40 (monomer, dimer, heterodimer), Alexa Fluor® 700 anti-mouse IL-12/IL-23 p40

## Product Data

---



IFN $\gamma$ -stimulated (two hours) Balb/c macrophages in the presence of LPS and monensin (24 hours) were fixed, permeabilized, and intracellularly stained with IL-12/IL-23 p40 (clone C15.6) PerCP/Cyanine5.5 (filled histogram) or rat IgG1, ? PerCP/Cyanine5.5 isotype control (open histogram).

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587