

## Brilliant Violet 785™ anti-mouse IFN-γ Antibody

<b>Catalog# / Size</b>	505837 / 125 µL 505838 / 50 µg
<b>Clone</b>	XMG1.2
<b>Regulatory Status</b>	RUO
<b>Other Names</b>	Interferon-γ, Immune interferon, Type II interferon, T cell interferon, Macrophage-activating factor (MAF)
<b>Isotype</b>	Rat IgG1, κ
<b>Description</b>	IFN-γ is a potent multifunctional cytokine which is secreted primarily by activated NK cells and T cells. Originally characterized based on anti-viral activities, IFN-γ also exerts anti-proliferative, immunoregulatory, and proinflammatory activities. IFN-γ can upregulate MHC class I and II antigen expression by antigen-presenting cells.

### Product Details

---

<b>Verified Reactivity</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Host Species</b>	Rat
<b>Immunogen</b>	<i>E. coli</i> -expressed, recombinant mouse IFN-γ
<b>Formulation</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
<b>Preparation</b>	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 785™ under optimal conditions.
<b>Concentration</b>	µg size: 0.2 mg/mL µL size: lot-specific (to obtain lot-specific concentration, please enter the lot number in our <a href="#">Concentration and Expiration Lookup</a> or <a href="#">Certificate of Analysis</a> online tools.)
<b>Storage &amp; Handling</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. <b>Do not freeze.</b>
<b>Application</b>	<a href="#">ICFC - Quality tested</a>
<b>Recommended Usage</b>	<p>Each lot of this antibody is quality control tested by <a href="#">intracellular immunofluorescent staining with flow cytometric analysis</a>. For flow cytometric staining using the µL size, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood. For flow cytometric staining using the µg size, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.</p> <p>Brilliant Violet 785™ excites at 405 nm and emits at 785 nm. The bandpass filter 780/60 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. <b>Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.</b> Refer to your instrument manual or manufacturer for support. Brilliant Violet 785™ is a trademark of Sirigen Group Ltd.</p> <p><a href="#">Learn more about Brilliant Violet™.</a></p> <p>This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.</p>
<b>Excitation Laser</b>	Violet Laser (405 nm)
<b>Application Notes</b>	<b>ELISA<sup>1-4,11,14</sup> or ELISPOT<sup>5</sup> Detection:</b> The biotinylated XMG1.2 antibody is useful as a detection

antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with purified R4-6A2 antibody (Cat. No. 505702/505706) as the capture antibody and recombinant mouse IFN- $\gamma$  (Cat. No. 575309) as the standard.

**ELISA or ELISPOT Capture:** The purified XMG1.2 antibody is useful as a capture antibody for a sandwich ELISA or ELISPOT assay, when used in conjunction with biotinylated R4-6A2 antibody (Cat. No. 505704) as the detection antibody and recombinant mouse IFN- $\gamma$  (Cat. No. 575309) as the standard. The LEAF™ purified antibody is suggested for ELISPOT capture (Cat. No. 505812).  
**Flow Cytometry**<sup>7,8,12,13,16</sup>: The fluorochrome-labeled XMG1.2 antibody is useful for intracellular immunofluorescent staining and flow cytometric analysis to identify IFN- $\gamma$ -producing cells within mixed cell populations.

**Neutralization**<sup>1-3,9,10</sup>: The XMG1.2 antibody can neutralize the bioactivity of natural or recombinant IFN- $\gamma$ . The LEAF™ purified antibody (Endotoxin <0.1 EU/ $\mu$ g, Azide-Free, 0.2  $\mu$ m filtered) is recommended for neutralization of mouse IFN- $\gamma$  bioactivity *in vivo* and *in vitro* (Cat. No. 505812). For *in vivo* studies or highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 505834) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/ $\mu$ g).

**Additional reported applications (for the relevant formats) include:** Western blotting, immunohistochemical staining of frozen tissue sections<sup>6,22,23</sup>, and immunocytochemistry.

**Note:** For testing mouse IFN- $\gamma$  in serum, plasma or supernatant, BioLegend's ELISA Max™ Sets (Cat. No. 430801 to 430806) are specially developed and recommended.

## Application References

1. Abrams J, *et al.* 1992. *Immunol. Rev.* 127:5. (ELISA, Neut)
2. Sander B, *et al.* 1993. *J. Immunol. Meth.* 166:201. (ELISA, Neut)
3. Abrams J, *et al.* 1995. *Curr. Prot. Immunol.* John Wiley and Sons, New York. Unit 6.20. (ELISA, Neut)
4. Yang X, *et al.* 1993. *J. Immunoassay* 14:129. (ELISA)
5. Klinman D, *et al.* 1994. *Curr. Prot. Immunol.* John Wiley and Sons, New York. Unit 6.19. (ELISPOT)
6. Sander B, *et al.* 1991. *Immunol. Rev.* 119:65. (IHC)
7. Ferrick D, *et al.* 1995. *Nature* 373:255. (FC)
8. Ko SY, *et al.* 2005. *J. Immunol.* 175:3309. (FC) [PubMed](#)
9. Peterson KE, *et al.* 2000. *J. Virol.* 74:5363. (Neut)
10. DeKrey GK, *et al.* 1998. *Infect. Immun.* 66:827. (Neut)
11. Dzhagalov I, *et al.* 2007. *J. Immunol.* 178:2113. (ELISA)
12. Lawson BR, *et al.* 2007. *J. Immunol.* 178:5366. (FC)
13. Lee JW, *et al.* 2006. *Nature Immunol.* 8:181. (FC) [PubMed](#)
14. Xu G, *et al.* 2007. *J. Immunol.* 179:5358. (ELISA) [PubMed](#)
15. Montfort M, *et al.* 2004. *J. Immunol.* 173:4084. [PubMed](#)
16. Haring JS, *et al.* 2008. *J. Immunol.* 180:2855. (FC) [PubMed](#)
17. Jordan JM, *et al.* 2008. *Infect Immun.* 76:3717. [PubMed](#)
18. Tonkin DR, *et al.* 2008. *J. Immunol.* 181:4516. [PubMed](#)
19. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
20. Cui Y, *et al.* 2009. *Invest. Ophthalm. Vis. Sci.* 50:5811. (FC) [PubMed](#)
21. Mykkanen OT, *et al.* 2014. *PLoS One.* 9:114790. [PubMed](#)
22. Yokogawa M, *et al.* 2013. *Mol. Carcinog.* 52:760. (IHC)
23. Mottram PL, *et al.* 1998. *J Immunol.* 161:602. (IHC)

## Product Citations

1. Lee J, *et al.* 2007. *Nat Immunol.* 8:181. [PubMed](#)
2. Hayashi K, *et al.* 2020. *Nat Commun.* 4.832638889. [PubMed](#)
3. Prosser A, *et al.* 2021. *Cell Reports.* 35(7):109141. [PubMed](#)
4. Halim TYF *et al.* 2018. *Immunity.* 48(6):1195-1207. [PubMed](#)
5. Farsakoglu Y *et al.* 2019. *Cell reports.* 26(9):2307-2315. [PubMed](#)
6. Walker JA, *et al.* 2020. *Immunity.* 51(1):104-118. [PubMed](#)
7. Marro BS, *et al.* 2019. *Cell Rep.* 29:3293. [PubMed](#)
8. Trittel S, *et al.* 2019. *Sci Rep.* 9:16362. [PubMed](#)
9. Montfort M, *et al.* 2004. *J Immunol.* 173:4084. [PubMed](#)

## RRID

AB\_11219004 (BioLegend Cat. No. 505837)  
AB\_2629667 (BioLegend Cat. No. 505838)

## Antigen Details

---

<b>Structure</b>	Cytokine; dimer; 40-80 kD (Mammalian)
<b>Bioactivity</b>	Antiviral/antiparasitic activities; inhibits proliferation; enhances MHC class I and II expression on APCs
<b>Cell Sources</b>	CD8 <sup>+</sup> and CD4 <sup>+</sup> T cells, NK cells
<b>Cell Targets</b>	T cells, B cells, macrophages, NK cells, endothelial cells, fibroblasts
<b>Receptors</b>	IFN- $\gamma$ R $\alpha$ (CDw119) dimerized with IFN- $\gamma$ R $\beta$ (AF-1)
<b>Cell Type</b>	Tregs

<b>Biology Area</b>	Cell Biology, Immunology, Neuroinflammation, Neuroscience
<b>Molecular Family</b>	Cytokines/Chemokines
<b>Antigen References</b>	<ol style="list-style-type: none"> <li>1. Fitzgerald K, <i>et al.</i> Eds. 2001. The Cytokine FactsBook. Academic Press, San Diego.</li> <li>2. De Maeyer E, <i>et al.</i> 1992. <i>Curr. Opin. Immunol.</i> 4:321.</li> <li>3. Farrar M, <i>et al.</i> 1993. <i>Annu. Rev. Immunol.</i> 11:571.</li> <li>4. Gray P, <i>et al.</i> 1987. <i>Lymphokines</i> 13:151.</li> </ol>
<b>Regulation</b>	Upregulated by IL-2, FGF-basic, EGF; downregulated by 1- $\alpha$ -25-Dihydroxy vitamin D3, dexamethasone
<b>Gene ID</b>	<a href="#">15978</a>

## Related Protocols


[Intracellular Cytokine Staining Protocol - Video](#)

[Intracellular Flow Cytometry Staining Protocol](#)

## Other Formats

APC anti-mouse IFN- $\gamma$ , Biotin anti-mouse IFN- $\gamma$ , FITC anti-mouse IFN- $\gamma$ , PE anti-mouse IFN- $\gamma$ , Purified anti-mouse IFN- $\gamma$ , Alexa Fluor® 488 anti-mouse IFN- $\gamma$ , Alexa Fluor® 647 anti-mouse IFN- $\gamma$ , Pacific Blue™ anti-mouse IFN- $\gamma$ , PerCP/Cyanine5.5 anti-mouse IFN- $\gamma$ , PE/Cyanine7 anti-mouse IFN- $\gamma$ , Brilliant Violet 421™ anti-mouse IFN- $\gamma$ , Brilliant Violet 650™ anti-mouse IFN- $\gamma$ , Ultra-LEAF™ Purified anti-mouse IFN- $\gamma$ , Brilliant Violet 711™ anti-mouse IFN- $\gamma$ , Brilliant Violet 785™ anti-mouse IFN- $\gamma$ , Brilliant Violet 605™ anti-mouse IFN- $\gamma$ , Brilliant Violet 510™ anti-mouse IFN- $\gamma$ , Purified anti-mouse IFN- $\gamma$  (Maxpar® Ready), PE/Dazzle™ 594 anti-mouse IFN- $\gamma$ , Alexa Fluor® 700 anti-mouse IFN- $\gamma$ , APC/Cyanine7 anti-mouse IFN- $\gamma$ , GolnVivo™ Purified anti-mouse IFN- $\gamma$ , APC/Fire™ 750 anti-mouse IFN- $\gamma$ , Spark NIR™ 685 anti-mouse IFN- $\gamma$

## Product Data



PMA + Ionomycin-stimulated (6 hours) C57BL/6 mouse splenocytes (in the presence of monensin) were stained with CD3 PE, fixed, permeabilized, and then stained with IFN- $\gamma$  (clone XMG1.2) Brilliant Violet 785™.

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 8999 BioLegend Way, San Diego, CA 92121 [www.biolegend.com](http://www.biolegend.com)  
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587